

Temporal variations of S wave attenuation field structure in the region of Zaili Alatau ridge: possible correlation with solar activity and large earthquakes

Inna N. Sokolova¹, Yuri F. Kopnichev², Galina Y. Khachikyan³

¹ Geophysical Survey of the Russian Academy of Sciences

² Schmidt Institute of physics of the Earth of the Russian Academy of science

³ Institute of Seismology, Ministry of Emergency Situations, RK

sokolovain@gsras.ru

We consider investigation results of spatio-temporal variations of S wave attenuation field in the lithosphere of the North Tien Shan.

It was shown that essential temporal variations of the attenuation field are observed in the middle and lower crust. We studied variations of Q_s value for the period of 1985-2017 using data of stations TLG and MDO, situated near Almaty city. We have been analyzing recordings of explosions, conducted at the Kotur-Bulak quarry. It is essential, that lines cross large Zaili deep fault zone for the both stations.

We studied variations of Q_s values in the middle and lower crust using coda wave envelopes at frequency of ~ 1 Hz. Note, that the lowest Q_s values ($\sim 60-100$) are observed in 1990-1995 for the both stations. It is important that the largest earthquakes in the North Tien Shan region for the last 40 years occurred in this time interval: Baisorun one (1990, $M_w=6.4$) and Suusamyr one (1992, $M_w=7.3$). It's possible to suggest that especially sensitive areas exist within zones of large regional faults. These areas can react on changes of the stress conditions within the crust sections, connected with large earthquake preparation, even located at considerable distances from such areas. (In the case of the Suusamyr earthquake such distances were about 250 km). The data on the attenuation field variations suggest that supersensitivity of such areas can be explained by active migration of deep-seated fluids in the middle and lower crust, which makes easier due to existence of large deep fault zones.

We found that a tendency of Q_s growth in the years of the low solar activity exists in comparison with the years of high solar activity. This can testify to influence of solar activity variations on fluid migration in the earth's crust of the region under investigation.

A comparison of the attenuation data with the location of the seismic activity zones shows that, as a whole, high attenuation corresponds to Zaili and Kemin zones. We found the most dangerous strips of these zones, where the lowest values of effective Q_s values are observed. First of all, such areas are located to the south-west and south-east of station MDO.