

Effect of interplanetary conditions and auroral activity on intermittency regularities of *Pi3* irregular geomagnetic pulsationsNadezhda Kurazhkovskaya¹, **Boris Klain**¹, Alexander Kurazhkovskii¹¹ Borok Geophysical Observatory, Branch of the Schmidt Institute of Physics of the Earth, Russian Academy of Sciences, Borok (Yaroslavl oblast), Russiaklb314@mail.ru

Earlier we showed that the cumulative distribution function (CDF) of the burst amplitudes of *Pi3* irregular geomagnetic pulsations observed during the development of substorms in the night sector of the magnetosphere follows the power law $f(A) = A^{-\alpha}$, where A is the amplitude of the bursts, α is the exponent. We considered the exponent α , which characterizes the CDF slope, as a characteristic of the intermittency and state of the medium in which *Pi3* bursts are formed. In this work are presented the results of study of the effect of the B_z component direction of the interplanetary magnetic field (IMF), auroral activity and b plasma parameter (equal to the ratio of thermal pressure to magnetic pressure) reflecting solar wind turbulence on the intermittency regularities of the *Pi3* pulsation bursts. One-minute digital observations of the magnetic field from two observatories in the Arctic and Antarctic: Heiss Island (HIS, corrected geomagnetic coordinates Latitude 74.80°, Longitude 144.46°) and Mirny (MIR, corrected geomagnetic coordinates Latitude -76.93°, Longitude 122.92°) from the World Data Center for Solar-Terrestrial Physics (Moscow) (http://www.wdcb.ru/stp/data/geo_min.val/) were used. Hourly average parameters of the interplanetary medium and auroral activity (AE -index) were obtained from the OMNI 2 database (<http://omniweb.gsfc.nasa.gov/ow.html>). A total of 305 and 415 *Pi3* pulsation bursts were analyzed in HIS and MIR, respectively. Taking into account interplanetary conditions, all *Pi3* events at each of the observatories were divided into two clusters depending on the direction of the IMF B_z component: *Pi3* bursts observed at $B_z < 0$ and $B_z > 0$. In a similar way, all cases were divided into two clusters according to the value of the AE index: *Pi3* bursts observed at $AE < 500$ nT and $AE > 500$ nT. According to the value of the b parameter, the events were divided into *Pi3*, observed at $0 < \beta \leq 1.3$ (solar wind plasma is highly turbulized) and $\beta > 1.3$ (weakly turbulized). For each group, a separate study of the CDF character of the *Pi3* pulsation amplitudes was performed and estimates of the power-law exponent α were obtained. It has been revealed that the bursts of *Pi3* pulsations observed in HIS and MIR are excited in a highly turbulent medium (the value of the exponent $\alpha > 1$). Regardless of the parameters considered (B_z , AE and β), in MIR the value of exponent α is significantly larger than HIS. For example, at $B_z < 0$ and $B_z > 0$ the value of α in HIS is 1.50 and 2.47, and in MIR 3.04 and 3.26, respectively. Similarly, at $AE < 500$ nT and $AE > 500$ nT, the α exponent in HIS is 1.60 and 1.52, in MIR 2.81 and 2.63, respectively. The distributions of *Pi3* burst amplitudes as a function of the parameter β (at $0 < \beta \leq 1.3$ and $\beta > 1.3$) were also approximated by power functions with lower α exponents in HIS (1.44 and 1.65) compared to MIR (2.60 and 2.74). It should be noted that similar trends are observed in the behavior of the exponent α in the northern and southern hemispheres when dividing *Pi3* bursts into clusters according to the values of B_z , AE and β . So, the value of the exponent α in both HIS and MIR is greater at $B_z > 0$, $AE < 500$ nT and $\beta > 1.3$ than at $B_z < 0$, $AE > 500$ and $0 < \beta \leq 1.3$. Thus, from the obtained results it follows that the level of turbulence in the region of excitation of *Pi3* pulsations (presumably in the magnetotail) is significantly higher in the absence of disturbances in the solar wind plasma, relatively moderate auroral activity and weak solar wind turbulence.