Distance corrections to surface-wave magnitudes of Far East shallow earthquakes

Nataliya V. Petrova¹ , Irina P. Gabsatarova² , Anna D. Kurova²

 1 Unified Geophysical Service of the Russian Academy of Sciences

² Geophysical Survey of the Russian Academy of Sciences

npetrova@gsras.ru

Surface wave magnitude MS has a number of advantages over other magnitude types - more than 100-year period of its determination, the ability to calculate MS for earthquakes around the world, established regional and global relationships with other magnitude types, participation of MLH (analogue of MS) in macroseismic field equations for the regions of the former USSR, which made it possible to use MLH and MS as a reference magnitude when compiling earthquake catalogs for seismic zoning of the USSR and Russia territories.

However, MS has two significant disadvantages - dependence on depth, which the authors of [1] proposed to compensate for using depth corrections, and on distance, as shown in this work.

In most seismological centers, MS is determined using the "Prague formula" [2]:

$$MS = lg(A/T)_{max} + 1.66 \times lgr^{0} + 3.3,$$

where A is the displacement in micrometers, T is the period in seconds corresponding to the maximum velocity, r is the epicentral distance ($2^{\circ} < r < 160^{\circ}$).

(1)

After the IASPEI adopted "Prague formula" as a standard (in 1967), the adequacy of the calibration function (the last two terms in formula (1)) was repeatedly discussed in publications and at conferences, but, despite some changes to the standard procedures for determining MS, the formula remained the same.

We checked the compliance of the calibration curve used in the "Prague formula" with the attenuation of A/T values depending on distance for Far East earthquakes with $h \le 50$ km for 2013–2018, the *MS* magnitudes of which are presented in the Seismological Bulletin of GS RAS (ftp ://ftp.gsras.ru/pub/Teleseismic_bulletin/). The distance dependence of the deviations *dMS* of station magnitudes *MSst* from the network average *MSav* was analyzed. For epicentral distances $r=4-80^{\circ}$, a loglinear dependence is established:

$$dMS = MSst - MSav = 0.661 \times \lg(r^{0}) - 1.06, R = 0.62, N = 8738.$$
 (2)

At distances $r=17-160^{\circ}$ absolute values of dMS do not exceed the generally accepted error in determining MS (±0.25), and only at $r<17^{\circ}$ MSst are significantly lower than average magnitudes MSav. When analyzing macroseismic data from 34 earthquakes in Northern Eurasia, we also discovered an underestimation of ISC and MOS magnitudes MS of weak earthquakes (MS<4.5) compared to the magnitude required in the regional macroseismic field equations. This is probably due to the big contribution to MSav of underestimated MSst due to small epicentral distances. Such an underestimation can be compensated for either by introducing corrections to MS for distance according to eq.(2) or by refining the calibration function.

REFERENCES

1. Petrova N.V., Gabsatarova I.P. (2020). Depth corrections to surface-wave magnitudes for intermediate and deep earthquakes in the regions of North Eurasia. Journal of Seismology, V. 24, 203-219. doi: 10.1007/s10950-019-09900-8

2. Vaněk J., Zátopek A., Kárnik V., Kondorskaya N., Riznichenko Y., Savarenski E., Solov'ev S., Shebalin N. (1962). Standardization of magnitude scales. Izv. Acad. Sci. USSR, Geophys Ser., 153-158: 108-111 (in English translation).