

Local time distribution and activity dependence of extreme electron densities in the auroral D-region as an image of energy-dependent energetic electron precipitation

Viktor Sergeev , Nikita Stepanov¹ , Yasunobu Ogawa , Eugene Rozanov

¹ Арктический и антарктический научно-исследовательский институт

victor40sergeev@gmail.com

Although electron density N_e in the auroral ionospheric D-region is formed by solar radiation and energetic particle precipitation, the highest observed densities are definitely produced by intense precipitated fluxes of energetic electrons (EE, here 30-300keV) and solar flare protons. Therefore, studying the most intense N_e episodes may provide valuable information about spatial distribution, spectral changes, and drivers of intense EE precipitation, which have important space weather implications. By combining EISCAT-Tromso UHF observations in different modes made in 2001-2021 (with polar cap absorption events being excluded), we survey N_e occurrence distributions and study statistical properties of EE precipitation producing the highest 10%, 5%, and 1% of N_e values at the altitudes between 100 and 75 km (corresponding to EE energies of ~ 20 to ~ 300 keV), and found two different patterns of spatial distribution of the extreme ionization. The largest N_e values occur in the nightside-early morning local time sector in the upper D-region ($H > 85$ km) and in the morning-noon sector at low altitudes (75-85 km). This bimodal pattern could partly be contributed by the suppressed N_e in the dark ionosphere where negatively charged ions may appear in large amounts. However, by analyzing measurements in the twilight conditions at all MLTs together with published statistical patterns of EE precipitation at different energies, we got indications that a dominance of pre-noon maximum at low altitudes in the D-region reflects the enhanced precipitation of >100 keV electrons in this local time sector. Statistically, at all altitudes (especially below 80/90km) the appearance of the highest N_e values favors enhanced auroral activity (not necessarily storms as revealed by SymH index) and shows a strong preference for high solar wind velocity (>550 km/s). These properties copy the known driver properties of energetic electrons. We discuss the advantages of exploring N_e data at specific altitudes (against direct measurements of EE precipitation) to investigate statistically the effects of specific precipitated energy components and the evolution of EE energy spectra.