

Long-term variations in characteristics of upper neutral atmosphere and ionosphere from spectrometric and radio sounding measurements

Irina Medvedeva¹ , Konstantin Ratovsky¹

¹ Institute of Solar-Terrestrial Physics, Siberian Branch of the Russian Academy of Sciences

ivmed@iszf.irk.ru

We present the results of studying year-to-year variations in the characteristics of the upper neutral atmosphere and the ionosphere. The analysis is based on the experimental data from the instrumental complex of the Institute of Solar-Terrestrial Physics SB RAS. As atmospheric and ionospheric characteristics we used the mesopause temperature (T_m) and peak electron density ($NmF2$); the analyzed period is 2008-2020. The mesopause temperature was obtained from spectrometric measurements of the OH emission ((6-2), 834 nm, Tory (51.8°N, 103.1°E)). The peak electron density was derived from the Irkutsk DPS-4 Digisonde measurements (52.3°N, 104.3°E). We analyzed the annual mean T_m and yearly average values of $NmF2$, as well as yearly average values of day-to-day and intradiurnal variability in T_m and $NmF2$. The analysis involved data on solar and geomagnetic activity, as well as on variations in the Southern Oscillation Index (SOI). To interpret the year-to-year variations, we use multiple regressions of the ionospheric and atmospheric characteristics on the F10.7- and Ap-indices of solar and geomagnetic activity, respectively. For the atmospheric characteristics, we also used regressions on the SOI-index as a characteristic of the lower atmosphere activity. It was revealed, that the yearly average values of $NmF2$ are dominantly controlled by changes in the solar flux. The year-to-year variations in the $NmF2$ variability are mainly driven by changes in both solar and geomagnetic activity. The year-to-year variations in the mesopause temperature weakly correlate with changes in the indices of solar and geomagnetic activity. The yearly average values of T_m variability correlate with changes in the SOI-index: the day-to-day variability demonstrates a positive correlation with the SOI-index, while the intradiurnal variability shows a negative correlation with the SOI-index. A significant relationship between the year-to-year variations in the $NmF2$ variability and T_m variability was not revealed.

The research was funded by the Russian Science Foundation (project No. 22-17-00146).